

Name:	Date		Date
Communication			Ambulate with cane
Person-Centered Care			Bleeding precautions
Wash hands			Wrist restraint
Standard Precautions			Cervical precautions
Handle Linen			Hip precautions
Occupied bed			Sternal precautions
Unoccupied bed			Oxygen on/off
Don and doff gloves			Seated transfer
Ambulate with gait belt			Thicken liquids
Amb. With gait belt and walker			Fracture pan
Position in bed			Regular bedpan
Position in chair			Urinal
Transfer bed to wheelchair			Change drainage bag to leg bag
Transfer wheelchair to bed			Change leg bag to drainage bag
Hoyer lift			Clean ostomy site
Hydration			Change ostomy bag
Assist with eating			Empty ostomy bag
Assist with toilet			Give enema
Change disposable brief			Give bowel suppository
Barrier creams/ointments			Catheter care
Dress/undress			External urinary catheter--on
Bed bath			External urinary catheter--off
Shower			Eyeglasses
Denture care			Hearing aids
Fingernail care			Elastic stockings
Foot care			Pediculicides
Mouth care			Shampoo
Perineal care			Mouth care—comatose client
Skin care			Shave--electric
Range of Motion--lower			Shave—safety razor
Range of Motion--upper			Sequential compression device
Weight			Warm therapy
Intake			Cold therapy
Output			Incentive spirometer
Pain level			Height
Temperature			Apical pulse
Radial pulse			Electronic blood pressure
Respirations			Manual blood pressure--forearm
Blood pressure—upper arm			Manual blood pressure—lower leg
Pulse oximetry			Manual blood pressure--thigh
Cough/deep breathe			Orthostatic blood pressure
Neutropenic precautions			
Isolation gown			
Isolation mask			
Collect specimen--urine			
Collect specimen--sputum			
Collect specimen--stool			
Abdominal thrust			